arXiv:1304.7232v1 [math.GT] 26 Apr 2013

THE A-POLYNOMIAL AND HOLONOMY PERTURBATIONS

JIANFENG LIN

Abstract. Dunfield-Garoufalidis and Boyer-Zhang proved that the A-polynomial of a nontrivial knot in S^3 is nontrivial. In this paper, we use holonomy perturbations to prove the non-triviality of the A-polynomial for a nontrivial, null-homotopic knot in an irreducible 3-manifold. Also, we give a strong constraint on the A-polynomial of a knot in the 3-sphere.

1. Introduction

In [11], an algebraic curve D_N was associated to a compact 3-manifold N with a single torus boundary. We briefly review the construction: consider the variety $\chi(\pi_1(N))$ of characters of $SL(2,\mathbb{C})$ representations of $\pi_1(N)$. We can restrict a representation to the boundary group $\pi_1(\partial N)$, which gives us a map $r\colon \chi(\pi_1(N)) \to \chi(\pi_1(T^2))$. Choose a basis $B=\{\mathfrak{M},\mathfrak{L}\}$ for the boundary group. Given two nonzero complex numbers m, l, we can define a $SL(2,\mathbb{C})$ representation $\gamma_{(m,l)}$ of $\pi_1(T^2)$:

$$\gamma_{(m,l)}(\mathfrak{M}) = \begin{pmatrix} m & 0 \\ 0 & m^{-1} \end{pmatrix}, \gamma_{(m,l)}(\mathfrak{L}) = \begin{pmatrix} l & 0 \\ 0 & l^{-1} \end{pmatrix} \tag{1}$$

So we have a natural map $h : \mathbb{C}^* \times \mathbb{C}^* \to \chi(\pi_1(T^2))$ defined by $h(m,l) := [\gamma_{(m,l)}] \in \chi(\pi_1(T^2))$.

Now we consider $h^{-1}(\bar{\mathrm{Im}}(r))$. The following lemma is proved in [3].

Lemma 1.1. *This algebraic set* $h^{-1}(\bar{Im}(r)) \subset \mathbb{C}^* \times \mathbb{C}^*$ *has no 2-dimensional irreducible components.*

The algebraic curve D_N is defined to be the union of the 1-dimensional components of $h^{-1}(\bar{\mathrm{Im}}(r))$. Each irreducible component of D_N is defined by a two variable irreducible polynomial. We multiply all these polynomials to get the A-polynomial of N, which we denote by $A_{N,B}$. (It depends on the choice of the basis B.)

If K is a null-homologous knot in a three manifold Y, we can take N to be the knot complement. There is a natural basis $\widetilde{B}$ for the boundary group: the meridian and the longitude (see Fact 2.7). The A-polynomial A_K of the knot K is defined to be $A_{N,\widetilde{B}}$.

In [3] and [12], Dunfield-Garoufalidis and Boyer-Zhang independently proved that:

Theorem 1.2 (Dunfield-Garoufalidis, Boyer-Zhang [3],[12])**.** *For a non-trivial knot* K *in* S^3*, we have* $A_K \neq l-1$.

Remark 1.3. *The reducible representations of the knot group always give an factor* $l-1$ *of the A-polynomial. The A-polynomial of a trivial knot is exactly* $l-1$*. Therefore, we say a knot* K *has non-trivial A-polynomial if* $A_K \neq l-1$*.*

Both Dunfield-Garoufalidis's and Boyer-Zhang's proofs make use of the following theorem.

Theorem 1.4 (Kronheimer-Mrowka [5])**.** *. For a non-trivial knot* K *in* S^3*, let* Y_r *(*$r \in \mathbb{Q}$*) be the manifold obtained by doing* r*-surgery along* K*. If* $|r| \leq 2$*, then* $\pi_1(Y_r)$ *admits some* $SU(2)$ *representation with non-cyclic image.*

This theorem is proved using techniques from $SU(2)$-gauge theory, namely holonomy perturbations. Holonomy perturbations were used by Floer in [9] to prove the surgery exact triangle. In fact, using holonomy perturbations, we can prove Theorem 1.2 directly. Moreover, we can prove something stronger.

The following are the main theorems of this paper.

Theorem 1.5. *Suppose* K *is a null-homologous, nontrivial knot in an orientable, closed, irreducible 3-manifold* Y*. If* $A_K = l-1$*, then there exists an homomorphism* $\rho : \pi_1(Y) \to SU(2)$ *such that* $\rho([K]) = -1 \in SU(2)$*.*

Theorem 1.6. *For a null-homotopic, nontrivial knot* K *in any orientable, closed, irreducible 3-manifold* Y*, we have* $A_K \neq l-1$ *.*

We have the following question.

Question 1.7. *Does any null-homologous knot in an irreducible 3-manifold have non-trivial A-polynomial?*

If we restrict to knots in S^3, Kronheimer and Mrowka's argument in [5] implies that the zero set of A-polynomial intersects the unit torus in some arcs. We give the concise statement:

Theorem 1.8. *Let* K *be a non-trivial knot in* S^3*. Then given any complex number* l *with* $|l| = 1$*, we can find another unit length complex number* m *such that* $A_K(l, m) = 0$*.*

Remark 1.9. *Theorem 1.8 gives strong constraints on the A-polynomial: for a generic polynomial, its zero set intersects the unit torus at some isolated points. This is ruled out by Theorem 1.8.*

In section 2, we review some preliminaries and basic constructions related to holonomy perturbations and non-vanishing theorem of critical points. In section 3, we prove the main theorems and give some examples.

Acknowledgement The author wishes to thank Nathan Dunfield, Hans Boden and Yi Ni for valuable discussions and comments. The author is especially grateful to Ciprian Manolescu for inspiring conversations and helpful suggestions in writing this paper.

2. PRELIMINARIES

In this section, we review some of the constructions related to holonomy perturbations. Most of the details can be found in [5] and [8].

2.1. **Holonomy perturbation.** M is a closed, oriented 3-manifold with $b_1(M) > 0$. Consider a rank 2 unitary bundle E over M with non-torsion $c_1(E)$. Let $\mathfrak{g}_E$ be the bundle whose sections are traceless, skew-hermitian endomorphisms of E. Let $\mathcal{A}$ be the affine space of $SO(3)$ connections of $\mathfrak{g}_E$. Let $\mathcal{G}$ be the group of gauge transformations on E with determinant 1.

Fix a reference connection A_0 on $\mathfrak{g}_E$. For any connection A on $\mathfrak{g}_E$, denoting $A - A_0$ by ω, we have the Chern-Simons functional:

$$CS : \mathcal{A} \to \mathbb{R}$$

$$CS(A) = \frac{1}{4}\int_{X_0} Tr(\omega \wedge d\omega + \frac{2}{3}\omega \wedge \omega \wedge \omega)$$

The critical points of the Chern-Simons functional are the flat connections.

Floer defined a class of perturbations of the Chern-Simons functional as follows. We take a function $\phi : SU(2) \to \mathbb{R}$ which is invariant under conjugation. It is uniquely determined by the even, 2π-periodic function $f(x) := \phi\begin{pmatrix} e^{ix} & 0 \\ 0 & e^{-ix}\end{pmatrix}$. Let D be a compact 2-manifold with boundary. Consider an embedding of $D \times S^1$ in M such that $\mathfrak{g}_E$ is trivial over it . Fix a trivialization of $\mathfrak{g}_E$ over $D \times S^1$ and take a 2-form μ which is supported in the interior of D and has integral 1. Using the trivialization, we can lift A to a connection $\bar{A}$ on the trivialized $SU(2)$ bundle over $D \times S^1$. Consider:

$$\Phi : \mathcal{A} \to \mathbb{R}$$

$$\Phi(A) = \int_{p\in D} \phi(Hol_{\{p\}\times S^1}(\bar{A}))\mu(p) \tag{2}$$

In this paper, we specialize D to be a disk D^2 or an annulus H. In the case of the disk, let $z_0 \in \partial D^2$ be a base point. We define the curves α and β to be $\partial D^2 \times \{0\}$ and $\{z_0\} \times S^1$ respectively. In the case of the annulus, we denote the two components of ∂H by c_1, c_2 and choose base points $z_i \in c_i$. Then for $i = 1, 2$, we define $\alpha_i, \beta_i \subset \partial(H \times S^1)$ to be $c_i \times \{0\}$ and $\{z_i\} \times S^1$ respectively.

Now we consider the perturbed Chern-Simons functional: $CS + \Phi : \mathcal{A} \to \mathbb{R}$.

The critical points can be completely described by the following lemmas. The first was proved by Floer. The second follows from similar arguments. For completeness, we sketch the proof for the second lemma. See [8] for details.

Lemma 2.1 (Floer [8])**.** *If $D \cong D^2$ and $A \in \mathcal{A}$ is a critical point of $CS + \Phi$, then:*
1) A is flat on $M - (D^2 \times S^1)$
2) After a suitable trivialization of $\mathfrak{g}_{D^2\times S^1}$, $Hol_\alpha(\bar{A}) = \begin{pmatrix} e^{i\tau} & 0 \\ 0 & e^{-i\tau}\end{pmatrix}$, $Hol_\beta(\bar{A}) = \begin{pmatrix} e^{i\nu} & 0 \\ 0 & e^{-i\nu}\end{pmatrix}$ and $\tau = f'(\nu) + 2\pi\mathbb{Z}$

Lemma 2.2. *If $D \cong H$ and $A \in \mathcal{A}$ is a critical point of $CS + \Phi$, then:*
1) A is flat on $M - (H \times S^1)$
2) The $SU(2)$ connection $\bar{A}$ is reducible and we can chose a suitable trivialization of $\mathfrak{g}|_{H\times S^1}$ such that:

$$Hol_{\alpha_1}(\bar{A}) = \begin{pmatrix} e^{i\tau_1} & 0 \\ 0 & e^{-i\tau_1} \end{pmatrix}, Hol_{\alpha_2}(\bar{A}) = \begin{pmatrix} e^{i\tau_2} & 0 \\ 0 & e^{-i\tau_2} \end{pmatrix}$$

$$Hol_{\beta_1}(\bar{A}) = Hol_{\beta_2}(\bar{A}) = \begin{pmatrix} e^{i\nu} & 0 \\ 0 & e^{-i\nu} \end{pmatrix}$$

Moreover, we have:

$$\tau_1 - \tau_2 = f'(\nu) + 2\pi\mathbb{Z} \tag{3}$$

Proof. Let P be the trivialized $SU(2)$ bundle over $H \times S^1$ and $\mathfrak{g}_P$ be the vector bundle associated to the adjoint representation. For $p \in H$, denote $\{p\} \times S^1$ by β_p. For a point $a = (p, q) \in H \times S^1$, we consider the holonomy of $\bar{A}$ along β_p. Let us denote it by $T_{\bar{A}}(a)$. This is an automorphism of the fiber of P over a. Thus $T_{\bar{A}}$ define a section of $Aut(P)$. It is easy to check that the gradient $(\nabla\Phi)(T_{\bar{A}})$ defines a section of $\mathfrak{g}_P$. Now A is a critical of $CS + \Phi$. It is proved in [8] that:

$$F_{\bar{A}} = \phi'(T_{\bar{A}})\mu \tag{4}$$

Using this, we can deduce that the holonomy along β_p does not depend on p and $\nabla_{\bar{A}}(T_{\bar{A}}) = 0$ (see [8]). If $T_{\bar{A}}(a) = \pm 1$ for any $a \in H \times S^1$, then $\phi'(T_{\bar{A}}) \equiv 0$. By equation (4), $\bar{A}$ is flat. Since $\pi_1(H \times S^1)$ is abelian, we see that $\bar{A}$ is reducible. Suppose $T_{\bar{A}}(a) \neq \pm 1$ for some $a \in H \times S^1$. Since $\nabla_{\bar{A}}(T_{\bar{A}}) = 0$, we have $T_{\bar{A}}(a) \neq \pm 1$ for any $a \in H \times S^1$. Thus the existence of this section tells us that $\bar{A}$ is also reducible in this case.

Because $\bar{A}$ is reducible, we can choose a suitable trivialization such that $\bar{A} = \sigma \cdot \left(\begin{smallmatrix} i & 0 \\ 0 & -i \end{smallmatrix}\right)$ for some one form $\sigma \in \Omega^1(H \times S^1)$. Since the holonomy along β_p does not depend on p, we can assume that $\mathrm{Hol}_{\beta_p}(\bar{A}) = \begin{pmatrix} e^{i\nu} & 0 \\ 0 & e^{-i\nu} \end{pmatrix}$ for any p. In particular, we have $\mathrm{Hol}_{\beta_i}(\bar{A}) = \begin{pmatrix} e^{i\nu} & 0 \\ 0 & e^{-i\nu} \end{pmatrix}$ for $i = 1, 2$. Then formula (4) becomes:

$$d\sigma = f'(\nu)\mu \tag{5}$$

By Stokes' theorem, we have $\int_{\alpha_1} \sigma - \int_{\alpha_2} \sigma = \int_{H \times \{0\}} d\sigma = f'(\nu)$. It is easy to see that $\tau_i = \int_{\alpha_i} \sigma$. Therefore, we proved the lemma. □

The following theorem was first proved in [5] and [7].

Theorem 2.3 (Kronheimer-Mrowka [5],[7])**.** *Let $M, E, \mathfrak{g}_E$ be defined as before. If M admits an oriented taut foliation and is not $S^2 \times S^1$, then for any holonomy perturbation Φ, the perturbed Chern-Simons functional $CS + \Phi$ over E has at least one critical point.*

This highly nontrivial theorem is a corollary of several theorems. We only sketch the proof here. For details, see [5] and [7]. If M admits taut foliation, then by the work by Eliashberg and Thurston in [13] and [14], M can be embedded in an admissible symplectic four manifold X. [5] proved that X can be chosen to satisfy several good conditions such that it satisfies Witten's conjecture relating Seiberg-Witten invariants and Donaldson invariants by the Feehan and Leness's work (See [7] and [9]). Because X is symplectic, its Seiberg-Witten invariants are non-trivial by Taubes's non-vanishing Theorem in [10]. From Witten's conjecture, it follows that the Donaldson invariants D_X^v are non-trivial for any line bundle v on X. Then by a standard stretching argument, the perturbed Chern-Simons functional has at least one critical point.

The following two theorems are proved by Gabai.

Theorem 2.4 (Gabai [2]). *Let M be a closed, orientable, irreducible 3-manifold. If $b_1(M) > 1$, then M admits a taut foliation.*

Theorem 2.5 (Gabai [2]). *Suppose $K \subset S^3$ is a non-trivial knot and Y_0 is obtained by doing 0-surgery along K. Then Y_0 admits orientable taut foliation and is not $S^2 \times S^1$.*

Combining these theorems we get:

Corollary 2.6. *If M is irreducible with $b_1(M) > 1$ or M is the 0-surgery manifold for a nontrivial knot in S^3, then for any holonomy perturbation, the perturbed Chern-Simons functional has a least one critical point.*

2.2. **The $SU(2)$-Pillowcase.** Now suppose $K \subset Y$ is a null-homologous knot in an orientable closed three manifold Y. Let $N(K)$ denotes the open tubular neighborhood of K. Unless otherwise stated, all homology groups are considered with $\mathbb{Z}$ coefficient. Using the Mayer-Vietoris sequence, it is easy to prove:

Fact 2.7. *The meridian $m \subset \partial(Y - N(K))$ is a primitive, nontorsion element in $H_1(Y - N(K))$. Also, we can find a unique longitude l such that $[l] = 0 \in H_1(Y - N(K))$ and $([m], [l])$ forms a basis for $H_1(\partial(Y - N(K)))$.*

Let $Y_r (r \in \mathbb{Q})$ be the manifold obtained by doing r-surgery along K. It is easy to see that $H_1(Y_0) = H_1(Y - N(K))$.

Suppose that the attached solid torus is $N_0 \subset Y_0$ with core c. Then $Y_0 - N_0 \cong Y - N(K)$. Now we consider a trivialized rank-2 unitary bundle E over $Y - N(K)$. Let $\mathfrak{g}_E$ be the bundle whose sections are traceless, skew-hermitian endomorphisms of E. Consider the flat connections A on $\mathfrak{g}_E$. Using the trivialization, we can lift A to a connection on the trivialized $SU(2)$ bundle $\tilde{P}$ over $Y - N(K)$. Denote this connection by $\tilde{A}$.

The following fact is standard:

Lemma 2.8. *By taking the holonomy of $\tilde{A}$, we get a 1-1 correspondence between flat connections on $\mathfrak{g}_E$ modulo gauge transformation and homomorphisms $\rho_A : \pi_1(Y - N(K)) \to SU(2)$ modulo conjugation.*

Because m and l are commutative as elements in $\pi_1(Y - N(K))$, we can do conjugation such that $\mathrm{Hol}_m(\tilde{A})$ and $\mathrm{Hol}_l(\tilde{A})$ are both diagonal.

Assume that

$$\mathrm{Hol}_\alpha(\tilde{A}) = \begin{pmatrix} e^{i\theta_A} & 0 \\ 0 & e^{-i\theta_A} \end{pmatrix}, \ \mathrm{Hol}_\beta(\tilde{A}) = \begin{pmatrix} e^{i\eta_A} & 0 \\ 0 & e^{-i\eta_A} \end{pmatrix}. \tag{6}$$

(There are some ambiguities here: (θ_A,η_A) are only well defined modulo involution $(a, b) \leftrightarrow (-a, -b)$ and translation $(a, b) \leftrightarrow (a, b) + 2\pi\mathbb{Z}^2$.)

Definition 2.9. *For a null-homologous knot $K \subset Y$, we define the subset of* $(\mathbb{R}/2\pi\mathbb{Z})^2$*:*

$$R_K := \{\pm(\theta_A, \eta_A) | A \textit{ is a flat connection on } \mathfrak{g}_E|_{Y_0 - N_0}\}$$

After checking the construction carefully and using Lemma 2.8, it's easy to see that R_K can be also defined as:

$$\{(\theta,\eta) | \exists \rho : \pi_1(Y-K) \to SU(2) \text{ s.t. } \rho(m) = \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix}; \rho(l) = \begin{pmatrix} e^{i\eta} & 0 \\ 0 & e^{-i\eta} \end{pmatrix}\} \quad (7)$$

Remark 2.10. *Because* $\pi_1(Y-K)$ *is finitely generated and* $SU(2)$ *is compact, the* $SU(2)$ *representation space of* $\pi_1(Y-K)$ *is compact. Thus* R_K *is a closed subset of* $(\mathbb{R}/2\pi\mathbb{Z})^2$*.*

Remark 2.11. *If* ρ *is a reducible* $SU(2)$ *representation of* $\pi_1(Y-K)$*, then* $\rho(l)=1$*. Thus the points in* R_K *off the line* $\eta = 2k\pi$ *give irreducible representations.*

The case $Y \cong S^3$ of the following lemma is proved as lemma 13 in [5]. Although the general proof is essentially the same, we give it here for completeness.

Lemma 2.12. R_K *is invariant under the translation by* $(\pi, 0)$*.*

Proof. By Fact 2.7, there exists $H_1(Y-N(K)) \to \mathbb{Z}_2$ mapping m to -1 and mapping l to 1. Considering the composition of $\pi_1(Y-N(K)) \to H_1(Y-N(K)) \to \mathbb{Z}_2 \to SU(2)$, we get $\rho_0 : \pi_1(Y-N(K)) \to SU(2)$ with the image in the center. For $\rho : \pi_1(Y-N(K)) \to SU(2)$, we multiply it with ρ_0 to get another representation ρ'. We have $\rho(m) = -\rho'(m)$ and $\rho(l) = \rho'(l)$. Then we use formula (7) to prove the lemma. □

3. Proof of main theorem

3.1. **The general case.** $K \subset Y$ is a null-homologous, nontrivial knot in an irreducible 3-manifold. We take two copies of $Y - N(K)$ and denote them by $Y_j - N(K_j)$ $(j = 1, 2)$. We glue them onto a thickened torus $T \times [-1, 1]$ using the same gluing map to get a closed manifold M. M is just the double of the knot complement. It's easy to see that $b_1(M) > 0$ and the meridian is a non-torsion element in $H_1(M)$.

Remark 3.1. *We identified the torus* T *with* $\partial(Y-N(K))$*. Thus it make sense to talk about the meridian* $m \subset T$ *and the longitude* $l \subset T$ *by the Fact 2.7.*

Now we consider the rank-2 unitary E with $c_1(E)$ the Poincaré dual of $[m] \in H_1(M)$. We can get $\mathfrak{g}_E$ by the following procedures:

- Construct trivialized $SO(3)$ bundles over $Y_j - N(K_j)$ $(j = 1, 2)$ and $T \times [-1, 1]$. Denote them by $\mathfrak{g}_1$, $\mathfrak{g}_2$ and $\mathfrak{g}_0$ respectively.
- Choose a map $f : T^2 \to SO(3)$ such that $f_* : \pi_1(\partial N_0) \to \pi_1(SO(3)) \cong \mathbb{Z}_2$ satisfies:
$$f_*([m]) = [0], f_*([l]) = [1]. \quad (8)$$
- Glue $\mathfrak{g}_1$, $\mathfrak{g}_0$ along $T \times \{-1\}$ using the identity map and glue $\mathfrak{g}_0$,$\mathfrak{g}_2$ along $T \times \{+1\}$ using f.

In order to do holonomy perturbations, we embed $H_j \times S^1 (j = 1, 2)$ into M by i_1, i_2 as follows:

$i_1(H_1 \times S^1) = T \times [-1, 0]$ and $i_1(* \times S^1)$ is parallel to $l \subset T^2 \times [-1, 0]$, which means that we use the holonomy along the longitude to do perturbation Φ_1.

$i_2(H_2 \times S^1) = T \times [0, 1]$ and $i_2(* \times S^1)$ is parallel to $m \subset T^2 \times [0, 1]$, which means that we use the holonomy along the meridian to do perturbation Φ_2.

Now we consider the perturbed Chern-Simons functional, $CS+\Phi_1+\Phi_2:\mathcal{A}\to\mathbb{R}$.

Notice that we already have trivializations of $\mathfrak{g}_E$ when restricted to $Y_j-N(K_j)$ and $T^2\times[-1,1]$. Thus for a connection $A\in\mathcal{A}$, we can restrict A to these pieces and use the trivializations to lift it to trivialized $SU(2)$ bundles. We have six loops: $m\times\{j\}\subset T^2\times[-1,1]$ and $l\times\{j\}\subset T^2\times[-1,1], j=-1,0,1$. Denote them by m_j, l_j.

If A is a critical point of $CS+\Phi_1+\Phi_2$, denote by $\widetilde{A}$ the $SU(2)$-lift of $A|_{\mathfrak{g}_0}$.

Since we only do perturbations in the interior of $T\times[0,-1]$ and $T\times[0,1]$, we have $\widetilde{A}|_{T\times\{j\}}$ flat for $j=-1,0,1$. Taking the holonomy of $\widetilde{A}|_{T\times\{j\}}$, we get a representation $\rho_j:\pi_1(T^2)\to SU(2)$.

Notation 3.2. *For $(\theta,\eta)\in(\mathbb{R}/2\pi\mathbb{Z})^2$, we denote by $\rho_{(\theta,\eta)}$ the representation $\pi_1(T^2)\to SU(2)$ which maps m to $\begin{pmatrix} e^{i\theta} & 0\\ 0 & e^{-i\theta}\end{pmatrix}$ and l to $\begin{pmatrix} e^{i\eta} & 0\\ 0 & e^{-i\eta}\end{pmatrix}$. We write $\rho\sim\rho'$ if two representations ρ and ρ' differ by a conjugation. It is easy to see that $\rho_{(\theta,\eta)}\sim\rho_{(\theta',\eta')}$ if and only if $(\theta,\eta)=\pm(\theta',\eta')$.*

Applying Lemma 2.2 to Φ_1 with $\alpha_1=m_{-1},\alpha_2=m_0,\beta_1=l_{-1},\beta_2=l_0$, we see that $\rho_{-1}\sim\rho_{(\theta_{-1},\eta_{-1})}$ and $\rho_0\sim\rho_{(\theta_0,\eta_0)}$ with:

$$\eta_{-1}=\eta_0 \text{ and } \theta_{-1}-\theta_0=f_1'(\eta_0). \tag{9}$$

Applying lemma 2.2 to Φ_2 with $\alpha_1=l_0,\alpha_2=l_1,\beta_1=m_0,\beta_2=m_1$, we see that $\rho_0\sim\rho_{(\hat{\theta}_0,\hat{\eta}_0)}$ and $\rho_1\sim\rho_{(\theta_1,\eta_1)}$ with:

$$\hat{\theta}_0=\theta_1 \text{ and } \hat{\eta}_0-\eta_1=f_2'(\theta_1). \tag{10}$$

Remark 3.3. *Recall that we choose $\phi_1,\phi_2:SU(2)\to\mathbb{R}$ to define Φ_1 and Φ_2. They give us f_1, f_2.*

We can also use the trivialization of $\mathfrak{g}_j (j=1,2)$ to lift A to flat $SU(2)$ connections over $Y_j-N(K_j)$. Denote them by $\widetilde{A}_1$ and $\widetilde{A}_2$. Because we glue $\mathfrak{g}_1$, $\mathfrak{g}_0$ using identity map, we have:

$$\mathrm{Hol}_{m_{-1}}(\widetilde{A})=\mathrm{Hol}_{m_{-1}}(\widetilde{A}_1),\mathrm{Hol}_{l_{-1}}(\widetilde{A})=\mathrm{Hol}_{l_{-1}}(\widetilde{A}_1)$$

Since we glue $\mathfrak{g}_0$ and $\mathfrak{g}_2$ using f, their trivializations do not agree on their common boundary. Different trivializations of the $SO(3)$ bundle give different $SU(2)$ lifts of $A|_{T\times\{2\}}$. By formula (8), it is easy to check:

$$\mathrm{Hol}_{m_1}(\widetilde{A})=\mathrm{Hol}_{m_1}(\widetilde{A}_2),\mathrm{Hol}_{l_1}(\widetilde{A})=-\mathrm{Hol}_{l_1}(\widetilde{A}_2)$$

Notation 3.4. *Let $S\subset(\mathbb{R}/2\mathbb{Z}\pi)^2$ be a subset. If h is an function with period 2π, we denote the set $\{(\theta,\eta+h(\theta))|(\theta,\eta)\in S\}$ by $S+(*,h)$ and the set $\{(\theta+h(\eta),\eta)|(\theta,\eta)\in S\}$ by $S+(h,*)$. We also denote the set $\{(\theta+a,\eta+b)|(\theta,\eta)\in S\}$ by $S+(a,b)$ for constant a,b.*

By Definition 2.9 and the discussion above, we see that

$$(\theta_{-1},\eta_{-1})\in R_K,(\theta_1,\eta_1)\in R_K+(0,\pi). \tag{11}$$

By formula (9), we have $(\theta_0, \eta_0) \in R_K + (-f_1', *)$. Since $\rho_{(\theta_0,\eta_0)} \sim \rho_0 \sim \rho_{(\hat{\theta}_0,\hat{\eta}_0)}$, we have $(\hat{\theta}_0, \hat{\eta}_0) = \pm(\theta_0, \eta_0)$. Notice that $R_K + (-f_1', *)$ is symmetric under the reflection at $(0, 0)$. We have $(\hat{\theta}_0, \hat{\eta}_0) \in R_K + (-f_1', *)$. By formula (10), we see that $(\theta_1, \eta_1) \in R_K + (-f_1', *) + (*, -f_2')$. Thus (θ_1, η_1) is an intersection point of $R_K + (-f_1', *) + (*, -f_2')$ and $R_K + (0, \pi)$. We get the following important proposition.

Proposition 3.5. *Suppose that $R_K + (-f_1', *) + (*, -f_2')$ does not intersect $R_K + (0, \pi)$, then the perturbed Chern-Simons functional $CS + \Phi_1 + \Phi_2$ has no critical point.*

Now we are ready to prove Theorem 1.5.


η
$\theta = g_1(\eta)$
$\eta = \pi$
θ
$\eta = g_2(\theta)$
▲ : points in R_K
○ : points in $R_K + (g_1, *)$
● : points in $R_K + (g_1, *) + (*, g_2)$
□ : points in $R_K + (0, \pi)$

Figure 1.

Proof. Suppose K has trivial A-polynomial. Then $R_K \setminus \{\eta = 2k\pi\}$ has only finitely many points by Lemma 1.1. If there exists no representation $\pi_1(Y) \to SU(2)$ mapping K to -1, then $(0, \pi) \notin R_K$. By lemma 2.12 $(k\pi, \pm\pi) \notin R_K$. So, we can choose a odd, 2π-periodic function g_1 such that the curve $\{g_1(\eta) = \theta, 0 < \eta < 2\pi\}$ does not intersect R_K. Then $R_K + (g_1, *)$ does not intersect $\{\theta = k\pi, 0 < \eta < 2\pi\}$. This can be done because there are only finitely many points to move away form the lines $\{\theta = k\pi, 0 < \eta < 2\pi\}$ and none of the them have η component equal to π. We find another odd, 2π-periodic function g_2 such that $R_K + (g_1, *) + (*, g_2)$ does not intersect the translation $R_K + (0, \pi)$. This can be done because $R_K + (g_1, *)$ has only finitely many points outside the line $\theta = 2k\pi$ and none of them have θ component equal to $k\pi$. $R_K + (0, \pi)$ has only finitely many points off the line

$\theta = \pi$. We can move these discrete points away from the one dimensional components. As long as $|g_2| < \frac{\pi}{2}$, the one dimensional component of $R_K + (g_1, *) + (*, g_2)$ and that of $R_K + (0, \pi)$ do not intersect.

Then we find even, 2π-periodic functions f_1, f_2 such that $f_j' = -g_j$ and use them to define perturbations Φ_j. By Proposition 3.5, the perturbed Chern-Simons functional $CS + \Phi_1 + \Phi_2$ has no critical point.

If K is nontrivial and contained in a 3-ball in Y, then K has nontrivial A-polynomial by Theorem 1.2. If K is not contained in a 3-ball, $Y - N(K)$ is irreducible and the boundary is incompressible. Doubling an irreducible three manifold along its incompressible boundary, we get another irreducible manifold. Therefore, M is irreducible with $b_1(M) > 0$. By Corollary 2.6, the fact that $CS+\Phi_1+\Phi_2$ has at least one critical point gives a contradiction. □

Theorem 1.6 is a trivial corollary of Theorem 1.5. Here are some other corollaries.

Remark 3.6. *If K is null homotopic, then it is easy to see that R_K has no points on the line $\theta = 0, 0 < \eta < 2\pi$. Therefore, we just need the perturbation Φ_2 if we only want to prove Theorem 1.6.*

Corollary 3.7. *If K is a non-trivial, null-homologous knot in an irreducible, orientable 3-manifold M and $\pi_1(M)/(K)$ has no $SO(3)$ representation with non-cyclic image, then $A_K \neq l - 1$. In particular, if $\pi_1(M)$ has no non-cyclic $SO(3)$ representation, then any null-homologous, nontrivial knot has nontrivial A-polynomial.*

Proof. Suppose K has trivial A-polynomial, then there exists $\rho : \pi_1(M) \to SU(2)$ such that $\rho(K) = -1$. The representation ρ induces a $SO(3)$-representation of $\pi_1(M)$ which factors through $\pi_1(M)/(K)$. By our assumption, this representation has cyclic image and so does ρ. Thus ρ factors through $H_1(M)$, which is impossible because K is null homologous and $\rho(K) \neq 1$ □

Example 3.8. *If $K \subset M$ is a non-trivial knot in Poincaré homology sphere and $2[K] \neq e \in \pi_1(M)$, then $A_K \neq l - 1$.*

Example 3.9. *Let $K \subset Y_1$ be a non-trivial knot in Y_1, where Y_1 is the manifold obtained by doing 1-surgery along some knot K' in S^3. If K is homotopic in Y_1 to the meridian of $S^3 - K'$ then $A_K \neq l - 1$.*

The following example is given by Dunfield:

Example 3.10. *Let Y be the $\frac{37}{2}$-surgery manifold on (-2,3,7)-pretzel knot. Then $\pi_1(Y)$ admits no non-cyclic $SO(3)$ representation. Therefore any nontrivial null-homologous knot $K \subset Y$ has non-trivial A-polynomial.*

3.2. The S^3 case.

Now let's go back to S^3 and use the technique in [5] to prove Theorem 1.8. Let K be a non-trivial knot in S^3. By Remark 1.3, we can assume that $A_K(m, l) = (l - 1)f(m, l)$. By Theorem 1.2, we have $f(m, l) \neq 1$. Denote the algebraic curve defined by $f(m, l)$ in $\mathbb{C} \oplus \mathbb{C}$ by C.

Consider its intersection with the unit torus $T^2 = \{(m, l) \mid |m| = |l| = 1\}$. It is a close subset of the torus. The following fact is straight forward by Lemma 1.1 and formula 7.

Lemma 3.11. *If* $(\theta, \eta) \in R_K$ *and* $\eta \neq 2k\pi$*, then either* $(e^{i\theta}, e^{i\eta})$ *is a* 0*-dimensional component of* $h^{-1}(\overline{Im}(r))$ *or* $(e^{i\theta}, e^{i\eta}) \in C \cap T^2$*.*

Now we can prove Theorem 1.8:

Proof. Suppose for some unit length $l_0 = e^{i\eta_0}$ ($\eta_0 \in [0, 2\pi)$), we can't find $m_0 = e^{i\theta_0}$ such that $f(m_0, l_0) = 0$. Since $C \cap T^2$ is a compact set, there exists $\epsilon > 0$ such that this holds for any $\eta_0' \in [\eta_0 - \epsilon, \eta_0 - \epsilon]$. Thus we can assume that $\eta_0 \neq 0$ and the line $\{l = e^{i\eta_0}\} \subset \mathbb{C} \oplus \mathbb{C}$ does not pass through the 0-dimensional components of $h^{-1}(\overline{\mathrm{Im}}(r))$. By Lemma 3.11, we see that R_K does not intersect the line $\eta = \eta_0$. Thus the translation $R_K + (0, -\pi)$ (denote it by S_K) does not intersect the line $\eta = \eta_0 - \pi$. By Definition 2.9, S_K is symmetric under the reflection at the origin. Thus S_K does not intersect the line $\eta = \pi - \eta_0$. We assume that $\eta_0 \geq \pi$ for convenience and the other cases can be proved similarly. Now consider the following line segments:

- $c_1 = \{(\theta, \eta_0 - \pi) | \theta \in [0, \pi]\}$;
- $c_2 = \{(\theta, \pi - \eta_0) | \theta \in [-\pi, 0]\}$;
- $c_3 = \{(0, \eta) | \eta \in [\pi - \eta, \eta_0 - \pi]$;
- $c_4 = \{(\pi, \eta) | \eta \in [0, \eta_0 - \pi]$
- $c_5 = \{(-\pi, \eta) | \eta \in [\pi - \eta_0, 0]\}$



Figure 2.

By the discussion before, S_K does not intersect c_1 and c_2. Since S^3 is simply connected, S_K does not intersect c_3, c_4, c_5. We can joint all these segments to get a broken line from $(-\pi, 0)$ to $(\pi, 0)$ and passing through $(0, 0)$. By Remark 2.10, we can find a small neighborhood N of this broken line such that $S_K \cap N = \emptyset$. It is easy to see that we can find an odd, 2π-periodic function $g : [-\pi, \pi] \to (-\pi, \pi)$ such that the graph of g lies in N.

Thus S_K does not intersect the graph of g. We can find an even function $f : [-\pi, \pi] \to \mathbb{R}$ with period 2π satisfying $f' = g$.

We consider the rank-2 unitary bundle E over Y_0 (recall that Y_0 is the 0-surgery manifold along K) with $c_1(E)$ the Poincaré dual of the meridian. We do a holonomy perturbation along the surgery solid tours in Y_0 using the even function f. By Lemma 2.1, the critical points of the perturbed Chern-Simons functional correspond to the intersection of S_K with the graph of g, which is empty. (We have S_K instead of R_K here, because E has nontrivial first Chern class.) So the perturbed Chern-Simons functional has no critical points. This contradicts Corollary 2.6. $\square$

This theorem has the following corollary, which was also proved by Boden in [15]:

Corollary 3.12. *For any nontrivial knot $K \subset S^3$, $deg_M A(M, L) \neq 0$.*

References

[1] A.Floer, *Instanton homology and Dehn surgery.* The Floer memorial volume, Progr. Math., vol.133, Birkhauser, Basel, 1995,pp. 195-256 no.3,77-97.

[2] D.Gabai, *Foliation and the topology of 3-manifolds.* J. Differential Geom.26(1987), no.3, 479-536.

[3] Nathan M. Dunfield, Stavros Garoufalidis, *Non-triviality of the A-polynomial for knots in S^3.* Algebr. Geom. Topol. 4 (2004), 1145C1153.

[4] Y.Ni, *Link Floer homology detects the Thurston norm.* Geom. Topol. 13, No.5, 2991-3019 (2009).

[5] P.B.Kronheimer, T.S.Mrowka, *Dehn surgery, the fundamental group and SU(2).* Math. Res. Lett. 11 (2004), no. 5-6, 741C754.

[6] P.B.Kronheimer, T.S.Mrowka, *Embedded surfaces and the structure of Donaldson's polynomial invariants*, J. Differential Geom. 41(1995), no.3,573-734.

[7] P.B.Kronheimer, T.S.Mrowka, *Witten's conjeture and Property P*, Geom. Topol. 8(2004) 295-310.

[8] P.J.Braam and S.K.Donaldson, *Floer's work on instanton homology, knots and surgery.* The Floer memorial volume, Progr. Math., vol.133, Birkhauser, Basel, 1995,pp. 195-256

[9] P.M.N.Feehan and T.G.Leness, *A general $SO(3)$-monopole corbordism formula relating Donaldson and Seiberg-Witten invariants.* Preprint, arXiv:math.DG/0203047,2003

[10] C.H.Taubes, *The Seiberg-Witten invariants and symplectic forms*, Math. Res. Lett.1(1994),no.6,809-822.

[11] D.Cooper, M.Culler, H.Gillet, D.Long and P.Shalen, *Plane curves associated to character varieties of 3-manifolds.* Invent. Math. 118(1994) 47-84.

[12] S. Boyer and X.Zhang, *Every nontrivial knot in S^3 has nontrivial A-polynomial.* Proc. Amer. Math. Soc. 133 (2005), 2813-2815.

[13] Y. Eliashberg, *Few remarks about symplectic filling.* Geom. Topol. 8 (2004), 277C293.

[14] Y. Eliashberg and W.Thurston, *Confoliations.* University Lecture Series, no. 13, American Mathematical Society, (1998).

[15] H. Boden, *Nontriviality of the M-degree of the A-polynomial*,to appear in Proceedings of the American Mathematical Society.

Jianfeng Lin
Department of Mathematics, University of California Los Angeles, Los Angeles, US
juliuslin@math.ucla.edu